

The Birth of Molecules

In less than a trillionth of a second, atoms can collide, interact and give birth to molecules. With lasers and molecular beams, it is now possible to witness the motions of molecules as one substance changes to another

by Ahmed H. Zewail

In 1872 railroad magnate Leland Stanford wagered \$25,000 that a galloping horse, at some point in its stride, lifts all four hooves off the ground. To prove it, Stanford employed English photographer Eadweard Muybridge. After many attempts, Muybridge developed a camera shutter that opened and closed for only two thousandths of a second, enabling him to capture on film a horse flying through the air [see illustration at top right]. During the past century, all scientific disciplines from astrophysics to zoology have exploited high-speed photography to revolutionize understanding of animal and mechanical motions that are quicker than the eye can follow.

The time resolution, or shutter speed, needed to photograph the ultrafast motions of molecules is beyond any conventional scale. When a molecule breaks apart into fragments or when it combines with another to form a new molecule, the chemical bonds between atoms break or form in less than a trillionth of a second, or one picosecond. Scientists have hoped to observe molecular motions in real time and to witness the birth of molecules: the instant at which the fate of the molecular reaction is decided and the

final products are determined. Like Muybridge, they needed to develop an ultrafast shutter, but it had to work 10 billion times faster than the 19th-century model.



For the past decade our research group at the California Institute of Technology has been developing techniques to observe the dynamics of molecules in real time. From 1985 to 1987 we refined our system of advanced lasers and molecular beams to a point at which we can now record the motions of molecules as they form and break bonds. The reaction can be seen as it proceeds from reactants through transition states and finally to products—chemistry as it happens.

Because transition states exist for less than a trillionth of a second, the time resolution should be shorter—a few quadrillionths of a second, or a few femtoseconds (one is equal to 10^{-15} second). A femtosecond is a smaller unit of time than a tick of even the finest atomic clock. A femtosecond is to a second what a second is to 32 million years. Furthermore, whereas in one second light travels nearly 300,000 kilometers—almost the distance between the earth and the moon—in one femtosecond light travels .3 micron—about the diameter of the smallest bacterium.

Alchemists in ancient Egypt, Greece, Arabia and China did not know about

TIME RESOLUTION of high-speed photography has improved 10 billion times as the technology has evolved from filming movements of animals to capturing the birth of molecules. In one second, a horse gallops 10 meters, as shown in photographs (top) taken by Eadweard Muybridge in 1887. In five trillionths of a second, hydrogen iodide collides with carbon dioxide, creating carbon monoxide, hydroxide and iodine. The illustration (bottom) is based on quantitative observations made at Caltech.

the fundamental importance of this time scale to the change of one substance to another, although they certainly did recognize the art of the transmutation. Only in this century have chemists been able to use a variety of tools to understand the sci-



AHMED H. ZEWAIL is the Linus Pauling Professor of Chemical Physics at the California Institute of Technology and has been a faculty member there since 1976. He has pioneered ultrafast laser femtochemistry to investigate the real-time motion of molecules in reactions. He received his B.S. from Alexandria University in Egypt and his Ph.D. from the University of Pennsylvania. Among the awards he has received are the 1989 King Faisal International Prize, the Harrison-Howe Award and the John Simon Guggenheim Fellowship. Zewail is a member of the National Academy of Sciences. This article is dedicated "to the memory of my dear friend Dick Bernstein, a giant in the field of molecular-beam science."

ence of molecular reaction dynamics.

At a molecular level a reaction starts when two molecules—call them *A* and *B*—are brought close enough together so that they begin to interact. As the molecules move even closer, other molecular species, which are neither *A* nor *B*, are formed until a new, stable molecule, *C*, is created.

One of these transitory species represents the stage at which the reaction proceeds irreversibly to products. This stage, technically called the transition state, is here denoted as $A \cdots B$. Hence,



The converse process is also possible: when molecule *C* is energized, it can pass through a transition state to form molecules *A* and *B*.



In any chemical reaction the motions

of the electrons and nuclei of atoms determine how the molecules interact, and those interactions in turn create the forces that govern the reaction's dynamics. The molecular motions can be rigorously delineated by the laws of quantum mechanics. In many cases, Newtonian, or classical, mechanics can simply give the transfer of energy and momentum during the course of the reaction, just as effectively as it describes the collision between two objects. If investigators can determine how molecular motions change during the critical transition phase, we can understand how new chemical bonds form and old ones disappear.

In practice, chemists do not keep track of every possible motion of every electron and nucleus in a molecular system. Instead they have discovered clever ways to represent the state of a system. Each state possesses a certain

amount of energy. The potential energy of a molecule when the atoms are a certain distance apart can be represented as a point on a graph. When all the states and their corresponding potential energies are graphed, they form a surface that has mountains and valleys. Molecular systems will spontaneously move from high-energy states (mountains) to low-energy states (valleys), but they require energy to move from valleys to mountains. The bottom of a valley indicates a stable state; the slopes around the valley represent the region of transition states, the different configurations for the atoms in proximity. Chemists technically define a transition state as a saddle point on a potential-energy surface.

If a molecular system is composed of two atoms and if the system's potential energy depends only on the distance that separates the atoms, the



solve the motion. The detector was made sensitive enough to probe the cyanide molecule when it was free or in the process of separating from the iodine atom. The sensitivity was achieved by using probing laser pulses of different colors in accordance with the spectrum of the molecules. We were thus able to observe the spectral frames changing with time as the reaction progressed from an intact ICN molecule to an iodine atom and molecular fragment, cyanide. We enjoyed discussing the results late into the night.

In 1987 we reported on the probing of the reaction of ICN to iodine and cyanide with enough time resolution to be able to observe the transition state $\{I \cdots CN\}$ directly. We obtained the real-time dynamics of the reaction and then deduced detailed features of the potential-energy surface [see illustration on opposite page].

That experiment made it possible to clock the breaking of an elementary chemical bond on a femtosecond time scale. The findings generated enthusiasm in the scientific community and inspired many more theoretical and experimental studies. The science writer Isaac Asimov provided an interesting description of the experiment. He compared the experiment with sticking a pin in a balloon only four billionths of an inch across—the size of an average molecule.

While we were still building the new femtosecond laser facility in 1986, Bernstein visited us at Caltech, and we

planned to probe the collision and bonding of two molecules, that is, a bimolecular reaction. At first we could not find a way to detect the start of these reactions: the time zero. Although bimolecular reactions last less than a picosecond, the two reactant molecules must first travel to meet each other—a journey that lasts about a million times longer than the reaction itself.

We found a solution to this problem when we studied the investigations of our colleagues at Orsay, France, and at the University of Southern California. By anchoring the two reagents together with a weak van der Waals bond, we could avoid the long time journey and establish the time zero. Fortunately, we had guidance from investigators at U.S.C. who had performed time-integrated studies on the reaction of hydrogen bromide and carbon dioxide.

For our first real-time experiments on these types of bimolecular reactions, we combined hydrogen iodide with carbon dioxide to produce iodine, carbon monoxide and hydroxide:



We wanted to understand this reaction at the elementary level. We broke the bond between hydrogen and iodine with a pump pulse. We then observed with probe pulses that the hydrogen attacks the carbon dioxide and strips the oxygen atoms from the carbon dioxide. The hydrogen atom then strips the oxygen atoms from the carbon dioxide.

dioxide molecule finally emerges five picoseconds after the start of the reaction, as we reported in 1987 and earlier this year.


The technique also allowed us to view the dynamics of the transitory collision complex HOCO in real time. We were able to relate this complex to the potential-energy surface representing hydrogen's approach to carbon dioxide. We are still examining the many subtle differences between our results and theory.

Just as physicists like to deal with the hydrogen atom for simplicity, chemical physicists like to understand elementary molecular systems. The simplest of all chemical reactions is that involving two atoms. I thought it would be interesting to direct some efforts at unraveling the dynamics of salt molecules such as sodium iodide:



Alkali halide reactions were the prototype for the "alkali age" of molecular-beam experiments, and it is insightful to study them in the "femtosecond age." Furthermore, researchers at the University of Toronto had investigated the emission spectra of the sodium iodide molecule during its transition to atoms of sodium and iodine.

What intrigued us about sodium iodide is a rather interesting potential-energy surface for the interaction of



FEMTOSECOND MOTIONS of the salt sodium iodide (NaI) reveal the molecular dynamics of the chemical bond linking sodium and iodine. In the potential-energy diagram at the left, the ionic (attractive) curve intersects the covalent (repulsive) curve. As a result, sodium iodide can break up into sodium and iodine, acting in a covalent manner, or sodium iodide can

exist in a high-energy bound state, alternating between covalent and ionic behavior. If sodium iodide dissociates into atoms, then the intensity of part of the spectrum will jump in steps, as shown in the experimental results (top at right). If sodium iodide is in its high-energy bound state, then the intensity of part of the spectrum will oscillate (bottom at right).

sodium and iodine atoms. If these atoms are brought together, they repel one another, particularly at short distances. On the other hand, if the same elements are brought together as oppositely charged ions of sodium and iodine, they attract each other. In fact, at a short enough distance, a stable salt composed of sodium and iodine ions will be formed. When the sodium and iodine form a covalent bond, they share electrons to create a stable energy state. When the atoms form an ionic bond, the sodium atom donates an electron to the iodine atom to create charged ions that attract.

Nature, however, does not work exactly in this way. Actually, the atoms behave as if they are both ionic and covalent. The true potential-energy curves are therefore composed of an ionic and a covalent curve [see illustration on preceding page]. The ionic curve crosses the covalent curve at a certain distance. In this region of interaction, the molecular system has a mixture of covalent and ionic characteristics; in other words, the molecule has a certain probability of being covalent and a certain probability of being ionic.

Theorists have divided the potential-energy curves into two parts: a lower and an upper. The lower curve, which represents low-energy states, is composed of the ionic curve at short distances and the covalent curve at long distances. The upper curve depicts the opposite situation: it is covalent at short distances and ionic at long distances. The upper curve also represents a state high in energy.

The behavior represented by the lower surface usually dominates when the sodium and iodine atoms are brought together slowly so that they begin to interact. In this case, an electron passes from sodium to iodine to create a stable molecule of sodium iodide. If, however, the atoms come together too rapidly, they can jump, with a certain probability, from the lower to the upper surface. The upper surface no longer represents the covalent repulsive (or slightly bound) curve and instead has a point of least energy (a minimum). Molecules in this potential minimum will thus be somewhat stable. (More specifically, the molecules enter a quasi-bound state.)

We hoped to view the femtosecond dynamics of the bond in sodium iodide as it breaks to form atoms of sodium and iodine. In particular, we focused on the jump from the upper potential surface to the lower one, or vice versa. We observed the motion as the molecules change from being covalent to being ionic and as they go through

the crossing region. The sodium atom gave an electron to the iodine atom at a distance of .7 nanometer. The sodium atom had, in effect, employed its electron as a harpoon to reel in the iodine atom.

To perform the experiment, we first sent a laser pulse to excite the NaI molecules. When the sodium separates about .25 nanometer from the iodine, the bond begins to break. A second pulse is then released to probe the reaction at a point when the sodium iodide bond is completely broken and a sodium atom and an iodine atom are formed. The probe pulse excites the sodium atom and causes it to emit yellow light. By probing the motion from the moment the bond breaks—time zero—until the birth of free sodium atoms, we can count, in real time, the number of sodium atoms that have appeared on the lower curve. But, more important, by detecting the quasi-bound transition species, $\text{Na}\cdots\text{I}$, which has different spectral properties from those of free sodium, we can observe the transition from $\text{Na}\cdots\text{I}$ to free sodium and iodine atoms.

Some of the sodium iodide molecules on the upper curve, when reaching the crossing area, will jump to the lower curve, and their bond will break to form sodium and iodine atoms. The molecules that do not jump but instead remain on the upper curve do not break a bond. They continue to vibrate until they do jump. Hence, pulses of sodium atoms should be evident only after each round-trip in the well of the upper curve. This phenomenon is exactly what we have observed. The results show the motion of the molecule during the breaking of a chemical bond and give the details of the potential energy governing the motion of sodium and iodine atoms.

Laboratories in the U.S. (at IBM) and in Germany (at Freiburg University) are now applying femtosecond techniques to observe the dynamics of many different types of reactions in the gas phase. At Caltech, we are continuing to probe elementary reactions and to examine more complex reactions that involve multiple births (for example, $\text{ABA} \rightarrow \text{AB} + \text{A} \rightarrow \text{A} + \text{B} + \text{A}$). We hope to study the simplest of all reactions, the combination of atomic hydrogen with molecular hydrogen.

We are also exploring large molecular systems with the aim of answering questions about selective reactivity. For example, if a molecule with two identical bonds is energized, do the two bonds break simultaneously or consecutively? To address this general and important question, we studied the dis-

sociation of $\text{C}_2\text{I}_2\text{F}_4$ into iodine and C_2F_4 . (In C_2F_4 , a double bond forms between the carbon atoms.) From real-time studies we found that the reaction proceeds consecutively, despite the molecular equivalence of the two carbon-iodine bonds. Even more remarkable is the discrepancy in the time scale for bond breakage—the first bond breaks in less than half a picosecond, whereas the second bond takes 100 times longer to break!

The ability to view molecular dynamics also suggests new ways of controlling reactions. There are several theoretical schemes for achieving control, and already experimentalists are exploring some new possibilities. The prospect exists for fine-tuning the motion and reactivity of molecules. If successful in the coming decades, laser-customized chemistry may be developed.

Muybridge invented high-speed photography for the fun of it, and today the technique has gone far beyond what he or Stanford could imagine. Although one cannot predict the future, we are certain of the importance and beauty of understanding the science of molecules and their reactions. The door is open for much experimental and theoretical research and for unexpected findings. It is my hope that readers will share in the fun and excitement that we have experienced in learning about the basics of alchemy—the transmutation of one substance to another—by observing the ultrafast motions of molecules.

FURTHER READING

- STATE-TO-STATE REACTION DYNAMICS. Richard N. Zare and Richard B. Bernstein in *Physics Today*, Vol. 33, No. 11, pages 43-50; November, 1980.
- INVESTIGATION OF ULTRAFAST PHENOMENA. Charles V. Shank in *Science*, Vol. 233, No. 4770, pages 1276-1280; September 19, 1986.
- THE NOBEL PRIZE FOR CHEMISTRY in *Les Prix Nobel 1986*. Almquist & Wiksell International, 1987.
- REAL-TIME LASER FEMTOCHEMISTRY: VIEWING THE TRANSITION FROM REAGENTS TO PRODUCTS. Ahmed H. Zewail and Richard B. Bernstein in *Chemical and Engineering News*, Vol. 66, No. 45, pages 24-43; November 7, 1988.
- LASER FEMTOCHEMISTRY. Ahmed H. Zewail in *Science*, Vol. 242, No. 4886, pages 1645-1653; December 23, 1988.
- EXPOSING MOLECULAR MOTIONS. Ian W. M. Smith in *Nature*, Vol. 343, No. 6260, pages 691-692; February 22, 1990.
- ULTRAFAST MOLECULAR REACTION DYNAMICS. L. Khundkar and Ahmed H. Zewail in *Annual Review of Physical Chemistry*, Vol. 41, pages 15-60; 1990.